

| Code (ICAR Central Health Key) | Equivalent code (former version of the recording standard) | Technical term | Synonyms, explanations | Repeated recording (y=yes, n=no, A=animal, P=parity) | Localizations | Average disease duration (i.e. new case after ...) |
|--------------------------------|--|--|--|--|--|--|
| 1.10.05.07. | | Asymmetric claws (AC) | significant difference in width, height and/or length between outer and inner claw which cannot be balanced by trimming | n | FR,FL,HR,HL | |
| 1.10.06.09.02.01. | | Concave dorsal wall (CD) | concave shape of the dorsal wall | n | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI,HRO,HLI,HLO) | |
| 1.10.05.04. | | Corkscrew claw (CC) | any torsion of either the outer or inner claw. The dorsal edge of the wall deviates from a straight line | n | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI,HRO,HLI,HLO) | |
| 1.10.07.10 | | Digital dermatitis (DD) | dermatitis digitalis, Mortellaro disease, Strawberry disease, infection of the digital and/or interdigital skin with erosion, mostly painful ulcerations and/or chronic hyperkeratosis/proliferation | y | FR,FL,HR,HL | 4 months |
| 1.10.07.11. | | Interdigital / superficial dermatitis (ID) | all kind of mild dermatitis around the claws that is not classified as digital dermatitis | y | FR,FL,HR,HL | 4 months |
| 1.10.06.07. | | Double sole (DS) | underrun sole, two or more layers of under-run sole horn | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI,HRO,HLI,HLO) | 4 months |
| 1.10.07.09. | | Heel horn erosion (HHE) | erosio unguulae, slurry heel, erosion of the bulbs, in severe cases typically V-shaped, possibly extending to the corium | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI,HRO,HLI,HLO) | 4 months |
| 1.10.06.15. | | Horn fissure (HF) | crack in the claw wall | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI,HRO,HLI,HLO) | 4 months |
| 1.10.06.15.01.02. | | Axial horn fissure (HFA) | vertical (longitudinal) crack in the inner claw wall | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI,HRO,HLI,HLO) | 4 months |
| 1.10.06.15.02. | 1.10.06.03. | Horizontal horn fissure (HFH) | horizontal sandcrack / horizontal crack in the claw wall | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI,HRO,HLI,HLO) | 4 months |
| 1.10.06.15.01.01 | | Vertical horn fissure (HFV) | vertical sandcrack, vertical (longitudinal) crack in the outer or dorsal claw wall | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI,HRO,HLI,HLO) | 4 months |
| 1.10.06.10. | | Interdigital hyperplasia (IH) | corns, tyloma, interdigital fibroma, interdigital growth of fibrous tissue | y | FR,FL,HR,HL | 4 months |
| 1.10.07.08.03. | | Interdigital phlegmon (IP) | foul-in-the-foot, footrot, interdigital panaritium, inflammatory edema in the interdigital space, symmetric painful swelling of the foot commonly accompanied with odorous smell with sudden onset of lameness | y | FR,FL,HR,HL | 4 months |
| 1.10.05.06. | | Scissor claws (SC) | tip of toes crossing each other | n | FR,FL,HR,HL | |

| | | | | | | |
|----------------|-------------|--|---|---|--|----------|
| 1.10.06.16. | | Sole hemorrhage (SH) | sole bruising, diffused and/or circumscribed red or yellow discoloration of the sole and/or white line | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI, HRO,HLI,HLO) | 4 months |
| 1.10.06.16.01. | | Sole hemorrhage diffused form (SHD) | diffused light red to yellowish discoloration | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI, HRO,HLI,HLO) | 4 months |
| 1.10.06.16.02. | | Sole hemorrhage circumscribed form (SHC) | clear differentiation between discolored and normal colored horn | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI, HRO,HLI,HLO) | 4 months |
| 1.10.07.08. | | Swelling of coronet and/or bulb (SW) | digital phlegmon, panaritium, digital inflammatory edema, uni- or bilateral swelling of tissue above horn capsule which may be caused by different conditions | y | FR,FL,HR,HL | 4 months |
| 1.10.07.01. | | Ulcer (U) | pododermatitis circumscripta purulenta (circumscribed purulent inflammation of the corium of the claws), ulceration of the sole area specified according to localization (zones) such as bulb ulcer, sole ulcer, toe ulcer/necrosis | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI, HRO,HLI,HLO) | 4 months |
| 1.10.07.03. | | Sole ulcer (SU) | pododermatitis solearis circumscripta purulenta et necroticans (circumscribed purulent and necrotizing inflammation of the corium of the claws), penetration through the sole horn exposing fresh or necrotic corium | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI, HRO,HLI,HLO) | 4 months |
| 1.10.07.03.04. | | Bulb ulcer (BU) | pododermatitis pulvinalis circumscripta purulenta et necroticans (circumscribed purulent and necrotizing inflammation of the corium of the claws at the bulb), heel ulcer, ulcer located at the bulb | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI, HRO,HLI,HLO) | 4 months |
| 1.10.07.03.01. | 1.10.07.04. | Toe ulcer (TU) | pododermatitis solearis circumscripta purulenta et necroticans (circumscribed purulent and necrotizing inflammation of the corium of the claws at the toe), ulcer located at the toe | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI, HRO,HLI,HLO) | 4 months |
| 1.10.07.12.01. | | Toe necrosis (TN) | necrosis of the tip of the toe with affection of bone tissue | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI, HRO,HLI,HLO) | 4 months |
| 1.10.05.08. | | Thin sole (TS) | sole horn yields (feels spongy) when finger pressure is applied | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI, HRO,HLI,HLO) | 4 months |
| 1.10.12. | | White line disease (WL) | separation of the white line with or without purulent exudation | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI, HRO,HLI,HLO) | 4 months |
| 1.10.12.01. | 1.10.06.04. | White line fissure (WLF) | separation of the white line which remains after balancing both soles | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI, HRO,HLI,HLO) | 4 months |
| 1.10.12.02. | 1.10.07.06. | White line abscess (WLA) | pododermatitis purulenta (septica) circumscripta parietalis et abaxialis, necro-purulent inflammation of the corium | y | FR,FL,HR,HL (FRI,FRO,FLI,FLO,HRI, HRO,HLI,HLO) | 4 months |